

# **Late Neoproterozoic (Pan-African) reactivations in the Mesoproterozoic Karagwe-Ankole Belt (KAB) in Kivu (RDC), Rwanda and Burundi: chronological framework and paleostress field.**

Damien DELVAUX<sup>1</sup>, Max FERNANDEZ-ALONSO<sup>1</sup>, Stijn Dewaele<sup>1</sup>, Mohamed LAGHMOUCH<sup>1</sup>, Gloire B. GANZA<sup>2</sup>, Guy M. ILOMBE<sup>2</sup>, Christian B. KALIKONE<sup>2</sup>, Espoir B. MUGISHO<sup>2</sup>, Yogolelo M. MUKOKYA<sup>2</sup>, Toussaint MUSHAMALIRWA<sup>2</sup>, Charles NZOLANG<sup>2</sup>, Evelyne N. SAFARI<sup>2</sup> & Robert N. WAZI<sup>2</sup>

*1. Royal Museum for Central Africa, Tervuren, Belgium (damien.delvaux@africamuseum.be)*

*2. Université Officielle de Bukavu, Bukavu, R.D. Congo*

The Karagwe-Ankole Belt (KAB) in the Kivus of the Democratic Republic of the Congo (DRC), Rwanda and Burundi is generally considered as a Mesoproterozoic belt, with a geodynamic evolution governed by the 1375 Ma Kibaran magmatic event, the  $\pm 1.0$  Ga Rodinia and the  $\pm 550$  Ma Gondwana (Pan-African) amalgamation events (Tack et al., 2010). Here we focus on the  $\pm 550$  Ma Gondwana event, which has been generally underestimated in previous works. This last event was first detected by “anomalously” young ages interpreted as isotopic rehomogenisations of “Lufilian” age in the granites of Rwanda, including the  $\pm 980$  Ma Tin granites which have been frequently cataclased (Gerards and Ledent (1970). Similar ages were obtained also for Pb/Pb, Rb/Sr on microcline from pegmatites (Monteyne-Poulaert, 1962), U/Pb on hydrothermal muscovite (Walemba, 2001) and U/Pb on monazite in a tectonic breccia in Burundi (Brinckmann et al., 2001).

We compiled the existing chronological datasets for the Karagwe-Ankole belt from the Paleoproterozoic to the late Cenozoic rifting. It evidences a multistage evolution (magmatism, mineralization, sedimentation, metamorphism, ductile and brittle deformation) and, in particular, the importance of Pan-African reactivations. In parallel, recent work in the neighboring areas (e.g. in Tanzania, Uganda and Kenya) showed that East-Central Africa appears much more unstable than previously thought as a consequence of the east Gondwana amalgamation. Fritz et al. (2013) evidenced early Pan-African formation of the East African Orogen between 650 and 615 Ma due to the collision and closure of the Mozambique ocean. Further E-W convergence occurred between 590 and 570 Ma (Fritz et al., 2013; Saalman et al., 2016), together with interaction between the Tanzania Craton and the Bangweulu Block between 590 and 550 Ma (Boniface & Appel, 2018). Late Pan-African reactivations under E-W shortening were caused by the Tanzania–Congo / Dharwar Cratons convergence at about 530 Ma (Fritz et al. 2013).

Field observations at various sectors of the KAB belt evidence brittle faulting that appear unrelated to and younger than the previous “Kibaran” magmatic, deformation and mineralization events (1.375 Ga or  $\sim 1.0$  Ga), but contemporaneous with the deformation of the Neoproterozoic (Cryogenian-Ediacaran) Itombwe series. Paleostress inversion allowed to reconstruct an older and widely expressed tectonic stress field characterized by a general E-W horizontal compression in a strike-slip to thrust faulting regime, and a younger brittle reactivation under N-S horizontal compression. The first brittle event occurred in brittle-ductile conditions and is interpreted as related to the late Pan-African event that affected the entire belt. The second brittle event is restricted to a few area and whitens younger deformations. This finding opens important perspectives for a revision of the regional geological map and also for the better understanding of the mineral resources.

## References

- Boniface, N. & Appel, P., 2018. Stenian - Tonian and Ediacaran metamorphic imprints in the southern Paleoproterozoic Ubendian Belt, Tanzania: Constraints from in situ monazite ages. *Journal of African Earth Sciences*, 133, 25-35.
- Brinckmann J., Lehmann, B., Hein, U., Höndorf, A., Mussallam, K., Weiser, T. & Timm, F., 2001. La géologie et la minéralisation primaire de l'or de la chaîne Kibarienne, nord-ouest du Burundi, Afrique orientale. *Geologische Jahrbuch Reihe, D 101*: 3-195.
- Cahen, L., Gerards, M., Theunissen, K., Lavreau, J., Villeneuve, M. Frautschi, J.M., 1976. Réunion de travail des 1, 2, et 3 septembre 1975: La géologie des terrains précambriens voisins du fossé tectonique occidental, spécialement dans les régions cises et part et d'autre de la partie sud du lac Kivu et du nord du lac Tanganyika au Kivu, au Rwanda et au Burundi. *Mus. roy. Afr. centr., Tervuren (Belg.); Dept. Geol. Min., Rapp. ann. 1975*, 143/170.
- Fritz, H., Abdelsalam, M., Ali, K., Bingen, B., Collins, A., Fowler, A., Ghebreab, W., Hauzenberger, C., Johnson, P., Kusky, T., Macey, P., Muhongo, S., Stern, R. & Viola, G., 2013. Orogen styles in the East African orogen: a review of the Neoproterozoic to cambrian tectonic evolution. *Journal of African Earth Sciences*, 86, 65-106.
- Gerards, J. & Ledent, D., 1970. Grands traits de la géologie du Rwanda, différents types de roches granitiques et premières données sur l'âge de ces roches. *Annales de la Société géologique de Belgique*, 93; 477-489.
- Monteyne-Poulaert, G., Delwiche, R., Cahen, L. & Safiannikoff, A., 1962. Ages de minéralisations pegmatitiques et filoniennes du Kivu méridional (Congo oriental), Indications préliminaires sur les âges de phases pegmatitiques successives, *Bulletin de la Société belge de Géologie, Paléontologie et Hydrologie*, 71(2), 272-295.
- Saalmann, K., Manttari, I., Nyakecho, C. & Isabirye, E., 2016. Age, tectonic evolution and origin of the Aswa Shear Zone in Uganda: Activation of an oblique ramp during convergence in the East African Orogen *Journal of African Earth Sciences*, 117, 303-330
- Tack, L., Wingate, M.T.D., De Waele, B., Meert, J., Belousova, E., Griffin, A., Tahon, A. & Fernandez-Alonso, M., 2010. The 1375 Ma "Kibaran event" in Central Africa: prominent emplacement of bimodal magmatism under extensional regime. *Precambrian Research*, 180, 63-84.
- Walemba, K.M.A., 2001. Geology, geochemistry, and tectono-metallogenic evolution of Neoproterozoic Gold deposits in the Kadubu area, Kivu, Democratic Republic of Congo. Ph.D Thesis University of the Witwatersrand, Johannesburg, South Africa, 491p.